

A Practical Guide to First-Order Multiplet Analysis in ^1H NMR Spectroscopy

Thomas R. Hoye,* Paul R. Hanson,^{1a} and James R. Vyvyan^{1b}

Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455

Received March 9, 1994^o

The ability to deduce the proper set of coupling constant (J) values from a complex first-order multiplet in a ^1H NMR spectrum is an extremely important asset. This is particularly valuable to the task of assigning relative configurations among two or more stereocenters in a molecule. Most books and treatises that deal with coupling constant analysis address the less useful operation of generating splitting trees to create the line pattern from a given set of J values. Presented here are general and systematic protocols for the converse, i.e., for deducing the complete set of J values from the multiplet. Two analytical methods (A, systematic analysis of line spacings, and B, construction of what can be called inverted splitting trees) are presented first. A reasonably thorough and systematic set of graphical representations of common doublet of doublets (dd's), ddd's, and dddd's are then presented. These constitute a complementary method for identification of J 's through visual pattern recognition. These approaches are effective strategies for extraction of coupling constant values from even the most complex first-order multiplets.

Introduction

Proton NMR spectroscopy is, arguably, the most powerful tool for structure assignment in most classes of organic molecules. Increased access to spectral data acquired on higher-field NMR spectrometers means that more and more of the resonances in routine spectra are first-order. While it is true that most modern spectrometers (as well as an increasing number of desktop personal computers) have software routines capable of performing rapid simulation of multispin systems, this approach to the analysis of complex first-order multiplets is often cumbersome, time consuming, or less than convenient. Simulation of a given multiplet is largely an empirical process that requires an initial determination or estimate of several of the individual coupling constants. Thus, the use of these computer-aided algorithms for the generation of simple first-order multiplets is less attractive than the ability to deduce a correct set of coupling constants upon simple visual inspection of any of the typically encountered line patterns. Graphical representations of common multiplet patterns are, in principle, quite useful for comparative analysis of actual data, but nearly all the published work has focused on non-first-order multiplets.² The general application of such representations is somewhat limited for two reasons: second-order multiplets are less frequently encountered at higher fields, and if one's spin system does not have the precise $\Delta\nu/J$ value of the calculated spectrum, the multiplet in question may look significantly different from the published representation.

The emergence of routine multidimensional NMR spectroscopy has been accompanied by a decline in the learning, teaching, and practice of the important skill of assigning first-order multiplets by inspection. Determination of coupling constants is still the most valuable

general method for assigning relative configurations of stereogenic centers in molecules. While the power of various routine two-dimensional NMR experiments is unarguable, it comes with a price. Data collection for 2-D experiments is always more time-consuming than for the simpler 1-D experiment, and in most settings access to magnet time is finite. Since 1-D and 2-D methods often provide complementary structural information, it is important that chemists maintain expertise with both.

This paper describes two related methods (A and B) that allow one to identify individual coupling constants within even the most complex first-order multiplets likely to be encountered in ^1H NMR spectra. It also provides a set of graphical representations (C and Tables 1-11) for assisting in empirical, visual pattern recognition.

A first-order multiplet arises when no two of the spins within an interacting multispin system have $\delta\nu/J \leq \sim 6$, and it always contains a symmetrical distribution of line positions about the midpoint of the multiplet (i.e., the chemical shift).³ In first-order multiplets the distance between the outermost pair of peaks is the sum of each of the coupling constants (ΣJ 's), a fact that we frequently find useful in assigning or verifying, for example, the last J value for an incompletely resolved complex multiplet.⁴ Often "special relationships" exist among the sets of coupling constants. We define these as cases where one of the coupling constants is equal to some combination of sums and/or differences among the remaining coupling constants. So defined, these special relationships always serve to reduce the number of lines in the multiplet and simplify (or complicate, depending on one's experience and point of view) the observed pattern. Finally, note that it is often, but by no means always, possible to determine a given coupling constant from either of a pair of spin-coupled resonances.

Methods for Deducing Coupling Constant Values

A. Systematic Analysis of Line Spacings. The task of extracting the actual values of coupling constants

(3) Caution must be exercised, however, because many second-order patterns (e.g., AB, AA'BB', AA'XX', and ABX) are also symmetrical.

(4) Often the sum of the J 's for non-first-order multiplets is also the distance between the outside lines of the multiplet, but this relationship deteriorates by the appearance of additional lines as $\Delta\nu/J$ becomes smaller and smaller.

* Abstract published in *Advance ACS Abstracts*, June 15, 1994.

(1) (a) University of Minnesota Graduate School Fellow, 1991-92. (b) Hercules Fellow, 1993-94.

(2) (a) Jackman, L. M., Sternhell, S. In *International Series of Monographs in Organic Chemistry*; Barton, D. H. R., Doering, W., Eds.; Oxford: Pergamon Press, 1969; Vol. 5: Applications of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry, Chapters 2 and 4 and references therein. (b) Wiberg, K. B., Nist, B. J. *The Interpretation of NMR Spectra*; W. A. Benjamin: New York, 1962. (c) Becker, E. D. *High Resolution NMR: Theory and Chemical Applications*, Academic Press: New York, 1980 and references therein.

Chart 1. Association between Coupling Constants and Line Spacings^a within dd's and ddd's

	Association	Line Spacing	Description
for dd's	J_L	{1 to 3} ^a (= {2 to 4})	larger J
	J_S	{1 to 2} (= {3 to 4})	smaller J
	$\Sigma J's = J_L + J_S$	{1 to 4}	sum of J's
	$J_L - J_S$	{2 to 3}	difference of J's
for ddd's	J_L	{1 to 5} (= {2 to 6} = {3 to 7} = {4 to 8})	largest J
where	J_M	{1 to 3} (= {2 to 4} = {5 to 7} = {6 to 8})	medium J
$J_L \geq J_M + J_S$	J_S	{1 to 2} (= {3 to 4} = {5 to 6} = {7 to 8})	smallest J
	$\Sigma J's = J_L + J_M + J_S$	{1 to 8}	sum of J's
	$J_M + J_S$	{1 to 4} (= {5 to 8})	sum of smaller two J's
	$J_M - J_S$	{2 to 3} (= {6 to 7})	difference of smaller two J's
	$J_L + J_M$	{1 to 7} (= {2 to 8})	sum of larger two J's
	$J_L - J_M$	{3 to 5} (= {4 to 6})	difference of larger two J's
	$J_L + J_S$	{1 to 6} (= {3 to 8})	sum of largest and smallest J's
	$J_L - J_S$	{2 to 5} (= {4 to 7})	difference of largest and smallest J's
for ddd's	J_L	{1 to 4} (= {2 to 6} = {3 to 7} = {5 to 8})	largest J
where	J_M	{1 to 3} (= {2 to 5} = {4 to 7} = {6 to 8})	medium J
$J_L \leq J_M + J_S$	J_S	{1 to 2} (= {3 to 5} = {4 to 6} = {7 to 8})	smallest J
	$\Sigma J's = J_L + J_M + J_S$	{1 to 8}	sum of J's
	$J_M + J_S$	{1 to 5} (= {4 to 8})	sum of smaller two J's
	$J_M - J_S$	{2 to 3} (= {6 to 7})	difference of smaller two J's
	$J_L + J_M$	{1 to 7} (= {2 to 8})	sum of larger two J's
	$J_L - J_M$	{3 to 4} (= {5 to 6})	difference of larger two J's
	$J_L + J_S$	{1 to 6} (= {3 to 8})	sum of largest and smallest J's
	$J_L - J_S$	{2 to 4} (= {5 to 7})	difference of largest and smallest J's

^a {i to j} = the separation in hertz between lines i and j.




Figure 1. Examples of ddd's where $J_L \geq J_M + J_S$ (case i) and where $J_L \leq J_M + J_S$ (case ii).

from within a given multiplet is most obvious for simple doublet of doublets (dd's). If the lines of the multiplet are numbered sequentially from, say, left to right (cf. entry a in Table 1), two (of the six) pairs of line spacings are associated with the smaller J value. As also summarized in the top portion of Chart 1, two more pairs of line spacings are associated with the larger J , and the remaining two pairs represent, respectively, the sum of and difference between the large and small J 's. The

distance between lines i and j is denoted as { i to j } throughout the discussion.

It is important to recognize that even though various sets of lines can have the same spacing, not all of those sets represent a coupling constant; some are coincidental. This is often a point of confusion. For example the distances between lines 1 to 2, 2 to 3, and 3 to 4 in entry b in Table 1 are all identical even though only {1 to 2} and {3 to 4} represent an actual J ; {2 to 3} is the difference between the two J 's (and {1 to 4} is the sum of the two J 's).

The situation for doublet of doublet of doublets (ddd's, bottom portion of Chart 1) is somewhat more complex, but still readily decipherable. For this treatment it is useful to define the J 's as J_s , J_m , and J_l to correspond to the smallest, medium, and largest J 's of the ddd, respectively. Again, the lines are numbered sequentially from left to right. The relative line intensities are important. In the absence of special relationships, all lines are of equal intensity, and for a ddd there is a total of eight lines [cf. the example in case i) in Figure 1]. One frequently encounters multiplets that contain line superposition, which is always accompanied by differential relative line intensities and a reduction in the total number of lines [cf. the example in case ii) in Figure 1]. Under any circumstances the sum of the relative line intensities will always equal 8 for a ddd (4 for a dd, 16 for a dddd, etc.). Lines of relative intensity greater than one are assigned more than one line number [e.g., the

sequence 1-(2/3/4)-(5/6/7)-8 for a 1:3:3:1 apparent quartet (ddd with three equivalent J 's) or the example in case ii)]. Be aware that "leaning" within a given multiplet, arising from intermediate $\Delta\nu/J$ values for which first-orderedness still holds, will distort the relative intensities from perfect integer ratios.

For doublet of doublet of doublets two situations can arise: case i where $J_1 \geq J_m + J_s$ and case ii where $J_1 \leq J_m + J_s$. A typical example for each case is shown in Figure 1.⁵ With the lines now numbered as described above and with reference to the bottom portion of Chart 1, one can assign the values of J_s , J_m , and J_1 in each of these multiplets by measuring the appropriate line spacings. The distance between lines 1 and 2 (i.e., {1 to 2}) always corresponds to the smallest coupling constant (J_s) and {1 to 3} always corresponds to the next smallest coupling constant (J_m). However, J_1 corresponds to {1 to 5} for case i but to {1 to 4} for case ii. The task of identifying J_1 from within dddd's (or higher multiplets) by this strategy is considerably more difficult. However, removing the smallest coupling (J_s) from a dddd, thereby creating a simplified ddd, permits application of the above strategy. On the other hand, this simplification is the first step in creating what we call here an inverted splitting tree, a process that is generalized next.

B. Inverted Splitting Tree Generation. The process of deconvoluting a first-order multiplet more complex than a ddd by the method described in A is not straightforward. We now describe a systematic approach that is applicable to even the most complex first-order multiplets. This strategy amounts to generation of an inverted splitting tree. Many readers are familiar with the process of generating the appearance of a first-order multiplet from a given set of J values, and many texts present the creation of splitting trees from a single line by sequential branching (most easily done proceeding from the largest J to the smallest). However, the ability to do the converse, to deduce the proper individual J 's from a given complex multiplet, is the more valuable yet more difficult skill to attain.


The total number of lines and the relative line intensities within a given multiplet are important parameters. Recall that dd's, ddd's, and dddd's with no special relationships will consist of 4, 8, and 16 lines, respectively, all of equivalent intensity and that the presence of special relationships among the coupling constants both reduces the total number of lines and alters the relative line intensities. The sum of the line intensities, appropriately normalized, will be identical for every multiplet of a given class (i.e., 4 for dd's, 8 for ddd's, and 16 for dddd's).

A general protocol for deducing the individual J 's for a given multiplet (illustrated in Chart 2 specifically for the ten-line 1:2:1:1:3:3:1:1:2:1 dddd corresponding to entry e of Table 9) consists of the following:

Step i: As discussed earlier, the distance between lines 1 and 2 (or the, say, left-hand-most pair) always represents the smallest J value of the multiplet [cf., J_s in panel i) of Chart 2]. If their relative intensity is 1:1, then the smallest J is unique; if it is 1:2 (or 1:3, etc.), then there are two (or three, etc.) identical smallest J 's.

Step ii: Identify the full set of pairs of lines separated by this smallest J value. This is perhaps the most diffi-

Chart 2. Protocol for Generating an Inverted Splitting Tree: Identification of Individual Coupling Constants as Applied to the dddd from Entry e of Table 9



cult step in the process. A dddd will contain eight such pairs. Each pair will have a partner pair symmetrically arranged by reflection through the midpoint of the multiplet. Those pairs associated with lines of intensities > 1 will be partially or totally coincident with other pairs. Thus, the total number of pairs associated with any single line is equal to the relative intensity of that line. As seen in panel ii), this is manifested in the number of times the ends of interconnecting arcs intersect a given line position (or vertical tick). That is, lines of intensities 1, 2, and 3 in the multiplet in Chart 2 have the ends of one, two, or three arcs, respectively, terminating at that line position [cf. panel ii)].

Step iii: Identify the centers of each of the pairs created in step ii, which collectively represent a new, simplified pattern (a ddd) as indicated by the dots in panel iii) as well as the tick marks in panel iv) in Chart 2. Notice that this submultiplet (in this instance a seven-line 1:1:1:2:1:1:1 ddd) is the residual pattern that would remain after selective decoupling of the spin responsible for the smallest J in the original multiplet. The spacing between the first (or last) pair of dots in this simplified multiplet represents the next smallest J of the original

(5) Notice that the examples chosen to illustrate cases i and ii were somewhat arbitrarily chosen. Examples could easily have been selected in which the case i and case ii multiplets contained fewer than and exactly eight total lines, respectively, rather than the converse.

Table 1. dd's

entry	Multiplet Appearance	J_x J_{13}	J_y J_{12}	Σ J 's	Special Relationships
a		12	4	16	
b		8	4	12	
c		6	4	10	
d		5	4	9	
e		4	4	8	$J_{13} = J_{12}$
f		3	4	7	
g		2	4	6	
h		1	4	5	
i		0	4	4	

H $\delta = 0.61$; dd; $J = 8.3, 3.8$ Hz; entry b
 H $\delta = 0.44$; dd; $J = 3.8, 3.8$ Hz; entry e

Table 2. ddd's Where $J_y = J_x$ (app dt's)

entry	Multiplet Appearance	J_x J_{14}	$J_y = J_x$ $J_{12} = J_{12'}$	Σ J 's	Special Relationships
a		20	4=4	28	
b		12	4=4	20	
c		9	4=4	17	
d		8	4=4	16	$J_{14} = J_{12} + J_{12'}$
e		7	4=4	15	
f		6	4=4	14	
g		5	4=4	13	
h		4	4=4	12	$J_{14} = J_{12}$
i		3	4=4	11	
j		2	4=4	10	
k		1	4=4	9	
l		0	4=4	8	

H $\delta = 3.72$, ddd, $J = 11.3, 11.3, 2.5$ Hz
 entry k

multiplet [i.e., $J_{ms(\text{medium small})}$ in panel iv); incidentally note that $J_s = J_{ms}$ for the example in Chart 2].

Step iv: The centers of each of these new pairs [diamonds in panel iv] collectively represent a new, further simplified, four-line pattern (a dd). The distance between the first (or last) pair of diamonds in panel iv) as well as the tick marks in panel v) represents the third smallest coupling constant, $J_{ml(\text{medium large})}$.

Step v: Repeat as necessary until all J 's have been established. One simple check for internal consistency is to verify that the sum of the determined coupling constants (ΣJ 's) is equal to the distance between the two outermost lines of the multiplet.

C. Graphical Representations (Tables 1–11). An alternative strategy for analysis of first-order multiplets is through visual pattern recognition. Many will find this approach complementary or preferable to the more analytical methods discussed above. We have generated a series of tables that shows systematic sets of first-order multiplets for the most commonly encountered spin systems. Representations of dd's and ddd's as well as of the more complex doublet of doublet of doublet of doublets (5 spins, dddd's) are included. Within any one table a single coupling constant, arbitrarily named J_z , is varied from large to small (usually to 0 Hz) values; the remaining coupling constants (J_y, J_x, J_w) are invariant throughout any one table. As a consequence, within any one table, entry a consists of relatively widely spaced, identical halves of the multiplet that converge as the variable

J_z decreases until the two halves are superimposed in the limit where the $J_z = 0$. Intermediate entries correspond to partially merged situations in which the two halves have undergone one or more sequential crossings of the innermost lines.

For some of the Table entries, special relationships (vide supra) exist among the J 's. These relationships, along with the magnitudes of J_w, J_x, J_y, J_z , and the ΣJ 's, are listed in the right-hand columns of each table. Note that although a specific (but arbitrary) set of J values has been chosen for illustrating the trends within any one table, those trends hold, of course, for any set of J 's having similar relative magnitudes and special relationships.

For any given table the specific line numbers that correspond to the generic J_w, J_x, J_y , and J_z are noted below the generic column heading. These line numberings change from table to table. For example, J_z in Table 1 is J_{13} (i.e., |1 to 3|) while J_z in Table 2 is J_{14} .

Note that the line numberings used in Tables 1–11 do not follow the rigorous convention used in sections A and B. Instead, the multiplet in entry a is simply numbered from left to right disregarding relative line intensities. Thus, each line has a single number at the outset (entry a), the crossing phenomena just mentioned are more easily tracked, identification of J 's within a visually matched multiplet is facilitated, and, importantly, the line spacings corresponding to every coupling constant, including the variable J_z , remain the same for all entries in the table.

Table 3. ddd's Where $J_y \approx J_x$

entry	Multiplet Appearance	J_x J_{12}	J_y J_{13}	J_z J_{12}	Σ J 's	Special Relationships
a		9	4	3	16	
b		8	4	3	15	
(c)		7	4	3	14	$J_{15} = J_{13} + J_{12}$
d		6	4	3	13	
e		5	4	3	12	
f		4	4	3	11	$J_{15} = J_{13}$
g		3	4	3	10	$J_{15} = J_{12}$
h		2	4	3	9	
i		1	4	3	8	$J_{15} = J_{13} - J_{12}$
l		0	4	3	7	

Table 4. ddd's Where $J_y = 2J_x$

entry	Multiplet Appearance	J_x J_{12}	J_y J_{13}	J_z J_{12}	Σ J 's	Special Relationships
a		10	4	2	16	
b		8	4	2	14	
c		7	4	2	13	
(d)		6	4	2	12	$J_{15} = J_{13} + J_{12}$
e		5	4	2	11	
f		4	4	2	10	$J_{15} = J_{13}$
g		3	4	2	9	
h		2	4	2	8	$J_{15} = J_{12}$
i		1	4	2	7	
j		0	4	2	6	

Included in each table is one (or two) representative multiplet(s) arising from the circled (or boxed) proton(s) taken from ^1H NMR spectra of known molecules.⁶ Although many of these presentations are simulations⁷ of tabulated J values (since they are taken from "pre-electronic storage" spectra), we have confirmed that this is an accurate representation of the actual spectrum by simulating multiplets for which the actual spectra were available. The examples in Tables 1, 4, 5, 9(e), and 10 (g) are taken from "real" rather than simulated spectra. The examples in all of the tables were selected to illustrate cases where special relationships exist among the J 's. These are particularly instructive since first-order multiplets with no special relationships give a full complement of 4, 8, 16, ... lines of identical intensity that are relatively easy to identify and interpret.

Table 1 and Tables 2–5 contain all the possible permutations of relative J values for dd's and ddd's, respectively. In other words, every conceivable first-order dd and ddd pattern can be approximately matched to one of the entries in these tables. On the other hand, only a somewhat arbitrarily chosen representative set (vide infra) of all possible dddd's is illustrated in Tables 6–11.

Table 1 shows the familiar series of typical doublet of doublets. Notice that there are several relative line

Table 5. ddd's Where $J_y > J_x$

entry	Multiplet Appearance	J_x J_{12}	J_y J_{13}	J_z J_{12}	Σ J 's	Special Relationships
(a)		9	6	1	16	
b		8	6	1	15	
(c)		7	6	1	14	$J_{15} = J_{13} + J_{12}$
(d)		6	6	1	13	$J_{15} = J_{13}$
e		5	6	1	12	$J_{15} = J_{13} - J_{12}$
f		4	6	1	11	
g		3	6	1	10	
h		2	6	1	9	
i		1	6	1	8	$J_{15} = J_{12}$
j		0	6	1	7	

(6) Spectral data were collected on a variety of spectrometers ranging from 300 to 500 MHz. (a) Examples from Tables 1, 5, and 9, entry e: Vyvyan, J. R. Unpublished results. (b) Examples from Tables 2, 6, entry f, 7, and 8: North, J. T. Ph. D. Dissertation, University of Minnesota, 1990. (c) Examples from Tables 3, 9, entry g, and 10, entry c: Peck, D. R. Ph. D. Dissertation, University of Minnesota, 1984. (d) Example from Table 4: Koltun, D. O., Vyvyan, J. R. Unpublished results. (e) Examples from Tables 6, entry c, and 11: Dellaria, J. F. Ph. D. Dissertation, University of Minnesota, 1982. (f) Example from Chart 3: Renner, M. K., Priest, O. P. Unpublished results.

(7) Simulation was performed with the VNMR version 4.1 software package on a Varian VXR spectrometer using a spectrometer frequency of 500 MHz and a line width ranging from 0.5 to 0.7 Hz.

spacings that repeat themselves during the progression of $J_{x(13)}$ from large to small. For example, entries a and h are each a pair of narrowly spaced outer and a pair of widely spaced inner lines. Similarly, entries b and g are

Table 6. dddd's Where $J_y = J_x = J_w$ (app dq's)

entry	Multiplet Appearance	J_x J_{12}	$J_y = J_x = J_w$ $J_{12} = J_{12} = J_{12}$	Σ J 's	Special Relationships
a		12	2=2=2	18	
b		8	2=2=2	14	
c		7	2=2=2	13	
d		6	2=2=2	12 $J_{15} = J_{12} + J_{12} + J_{12}$	
e		5	2=2=2	11	
f		4	2=2=2	10 $J_{15} = J_{12} + J_{12}$	
g		3	2=2=2	9	
h		2	2=2=2	8 $J_{15} = J_{12} = J_{12} = J_{12}$	
i		1	2=2=2	7	
j		0	2=2=2	6	

$\delta = 2.23$, dddd, $J = 13.9, 4.3, 4.2, 4.0$ Hz
entry c

$\delta = 2.03$, dddd, $J = 13, 6.5, 6.5, 6.5$ Hz
entry f

Table 8. dddd's Where $J_y = J_x$ (app dtd's)

entry	Multiplet Appearance	J_x J_{17}	$J_y = J_x$ $J_{12} = J_{12}$	J_w J_{12}	Σ J 's	Special Relationships
a		14	4=4	2	24	
b		11	4=4	2	21	
c		10	4=4	2	20 $J_{17} = J_{15} + J_{15} + J_{15}$	
d		9	4=4	2	19	
e		8	4=4	2	18 $J_{17} = J_{15} + J_{15}$	
f		7	4=4	2	17	
g		6	4=4	2	16 $J_{17} = J_{15} + J_{12}$	
h		5	4=4	2	15	
i		4	4=4	2	14 $J_{17} = J_{15} = J_{12}$	
j		3	4=4	2	13	
k		2	4=4	2	12 $J_{17} = J_{12}$	
l		1	4=4	2	11	
m		0	4=4	2	10	

$\delta = 2.80$, dddd, $J = 12.0, 3.0, 3.0, 1.0$ Hz
entry a

Table 7. dddd's Where $J_x = J_y$ and $J_x = J_w$ (app tt's)

entry	Multiplet Appearance	$J_x = J_y$ $J_{14} = J_{14}$	$J_x = J_w$ $J_{12} = J_{12}$	Σ J 's	Special Relationships
a		8=8	2=2	20	
b		6=6	2=2	16	
c		5=5	2=2	14	
d		4=4	2=2	12 $J_{14} = J_{12} + J_{12}$	
e		3=3	2=2	10	
f		2=2	2=2	8 $J_{14} = J_{12}$	
g		1=1	2=2	6 $J_{12} = J_{14} + J_{14}$	
h		0=0	2=2	4	

$\delta = 3.95$, dddd, $J = 6.2, 6.2, 5.1, 5.1$ Hz
entry e

Table 9. dddd's Where $J_x = J_w$ (app dtd's)

entry	Multiplet Appearance	J_x J_{17}	J_y J_{12}	$J_x = J_w$ $J_{12} = J_{12}$	Σ J 's	Special Relationships
a		12	5	3=3	23	
b		11	5	3=3	22 $J_{17} = \Sigma J_{12,12,12}$	
c		10	5	3=3	21	
d		9	5	3=3	20	
e		8	5	3=3	19 $J_{17} = J_{15} + J_{12}$	
f		7	5	3=3	18	
g		6	5	3=3	17 $J_{17} = J_{12} + J_{12}$	

$\delta = 0.61$, dddd, $J = 8.7, 5.3, 3.4, 3.4$ Hz
entry e

$\delta = 1.96$, dddd, $J = 14.0, 8.2, 7.0, 7.0$ Hz
entry g

each a set of four equally spaced lines; they also have the same special relationship. Finally, the number of lines is reduced to three (cf. entry e) in the trivial and common case of a dd appearing as an apparent triplet with relative line intensities of 1:2:1.

Tables 2–5 show the multiplets associated with doublet of doublet of doublets under circumstances where two of the three coupling constants are equal ($J_{y(12')} = J_{x(12)}$, Table 2), nearly equal ($J_{y(13)} \approx J_{x(12)}$, Table 3), related by a factor of two ($J_{y(13)} = 2J_{x(12)}$, Table 4), or very different ($J_{y(13)} \gg J_{x(12)}$, Table 5). Table 2 contains the case where the two invariant coupling constants ($J_{y(12')}$ and $J_{x(12)}$) of

Table 10. dddd's Where $J_y \approx J_x \approx J_w$

entry	Multiplet Appearance	J_x J_{19}	J_y J_{14}	J_z J_{13}	J_w J_{12}	Σ J' 's	Special Relationships
a		12	5	4	3	24	$J_{19} = \Sigma J_{14, 12, 12}$
b		10	5	4	3	22	
c		9	5	4	3	21	$J_{19} = J_{14} + J_{12}$
d		8	5	4	3	20	$J_{19} = J_{14} + J_{12}$
e		7	5	4	3	19	$J_{19} = J_{12} + J_{12}$
f		6	5	4	3	18	$J_{19} = J_{14} + J_{12} - J_{12}$
g		5	5	4	3	17	$J_{19} = J_{14}$
h		4	5	4	3	16	$J_{19} = J_{12}$
i		3	5	4	3	15	$J_{19} = J_{12}$
j		2	5	4	3	14	$J_{19} = J_{14} - J_{12}$
k		1	5	4	3	13	$J_{19} = J_{12} - J_{12}$
l		0	5	4	3	12	

$\delta = 1.83$, dddd, $J = 14.0, 7.6, 6.5, 5.0$ Hz
entry c

$\delta = 4.23$, dddd, $J = 6.5, 6.5, 5.3, 3.8$ Hz
entry g

Table 11. dddd's Where $J_y \gg J_x \approx J_w$


entry	Multiplet Appearance	J_x J_{19}	J_y J_{15}	J_z J_{13}	J_w J_{12}	Σ J' 's	Special Relationships
a		12	7	3	2	24	$J_{19} = J_{15} + J_{13} + J_{12}$
b		11	7	3	2	23	
c		10	7	3	2	22	$J_{19} = J_{15} + J_{12}$
d		9	7	3	2	21	$J_{19} = J_{15} + J_{12}$
e		8	7	3	2	20	$J_{19} = J_{15} + J_{13} - J_{12}$
f		7	7	3	2	19	$J_{19} = J_{15}$
g		6	7	3	2	18	$J_{19} = J_{15} + J_{12} - J_{12}$
h		5	7	3	2	17	$J_{19} = J_{15} + J_{12}$

$\delta = 2.23$, dddd, $J = 15.5, 10.6, 5.1, 3.0$ Hz
entry c

$\delta = 1.41$, dddd, $J = 12.3, 9.4, 7.4, 3.0$ Hz
entry d

the ddd are identical so that the multiplets are all apparent doublets of triplets (6 lines), sometimes with additional simplification arising from additional special relationships [e.g., entries d (5 lines), h (4 lines), and l (3

Chart 3. Inverted Splitting Tree Analysis for the ddq (app tq) from H(3) in One Rotamer of the Mosher Amide 1



lines)]. Table 3 contains the cases where the two invariant coupling constants are only slightly different. The illustrated example, in which the central two lines of the ddd coincide and for which the largest J value is the sum of the two smaller J 's (i.e., $J_{x(15)} = J_{y(13)} + J_{z(12)}$ cf. entry c), is a commonly encountered one. This relationship and pattern (a seven-line 1:1:1:2:1:1:1 multiplet) are also seen in the example accompanying Table 4 (entry d). Table 5 illustrates the case where one of the invariant J 's is always significantly larger than the second ($J_{y(13)} \gg J_{z(12)}$). The multiplet arising from the boxed proton in the example molecule corresponds to a situation intermediate between entries c and d and illustrates an important point. Namely, to find the best fit one must sometimes envision the continuum of multiplets that arise by sliding together the two halves of the pattern for entry a (as J_z decreases) within any one table. There often is not a perfect correlation between the multiplet under analysis and a specific table entry.

Tables 6–11 show doublet of doublet of doublet of doublets (dddd's) that increase in their complexity as the tables progress. Table 6 illustrates the case where three of the four J 's are equal, resulting in an apparent doublet

of quartets (i.e., app dq). This type of relationship is prevalent in rigid ring systems either where a geminal and two diaxial couplings give three approximately equivalent J 's and the fourth and smallest J arises from the gauche axial-equatorial arrangement of a pair of vicinal protons or, as is the case for the example of the boxed proton (cf. entry c), where a single large geminal coupling is accompanied by three approximately equivalent and small gauche couplings. The special relationship where the largest J is twice as large as any of the equivalent smallest J 's is present in the example of the circled proton (cf. entry f) and results in the greatly simplified six-line pattern. Table 7 shows multiplets that contain two pairs of identical J 's (i.e., app tt's). The example in Table 7 (cf. entry e) is commonly encountered in acyclic molecules where a methine proton is flanked by two methylene groups that each constitutes a pair of diastereotopic protons. Table 8 depicts dddd's in which the two J 's of intermediate magnitude are identical (i.e., app dtd's). Notice that the smallest J (1 Hz) in the example multiplet (cf. entry a) arises from four-bond W-coupling. Table 9 presents multiplets in which the two smallest J 's are identical in magnitude (i.e., app ddt's), although cases where the value of $J_{z(17)}$ drops below $J_{y(13)}$ are not shown. Another commonly encountered ddt arises from the methine of a terminal allyl group (i.e., $\text{CH}_2=\text{CHCH}_2\text{R}$, not shown).

Tables 10 and 11 include representative cases of dddd's to which no universal special relationship applies (as is the case for Tables 6–9). Thus, for the first time among Tables 6–11 some of the entries show multiplets having the full complement of sixteen lines. The multiplets in Table 10 contain three and those in Table 11 contain two small J 's of nearly equal magnitude.

Recognize that there exist other possible combinations for the relative magnitudes of J 's within the family of dddd's than those shown in Tables 6–11. Recall that *all* permutations of J values for the simpler dd's and ddd's were encompassed by Tables 1 and 2–5, respectively. When in need, the reader is encouraged to extend the methods described here to those cases of J value combinations for dddd's not explicitly covered as well as to yet more complex multiplets like the ddddd's.

We conclude with one example of the latter. Chart 3 shows the multiplet arising from H(3) in the minor rotamer of the indicated Mosher amide.^{6f} The multiplet was identified by the inverted splitting tree method as a ddq with $J = 9.5, 9.5,$ and 6.5 by the sequence of steps outlined in Chart 3. This set of coupling constants requires that the molecule exists largely in the conformation having dihedral angles of $\sim 150^\circ$ and 30° between H(3)/H(4a) and H(3)/H(4b), respectively. It should be noted that it was not possible to identify the two vicinal 9.5 Hz J 's from the benzylic methylene resonances due to significant overlap among those resonances for the amide rotamers, even at 500 MHz.

We have found the complementary approaches of visual pattern recognition (C) as well as the more systematic and analytical methods described in A and B to be effective and powerful tools. These approaches permit easy access to the rich, intrinsic information that is universally available via the simplest of NMR experiments.

Acknowledgment. This work was supported by the DHHS Grants GM 34492 and 39339. We acknowledge Mr. Zhixiong Ye for enunciating ideas that aided the development of the systematic analysis of line spacings and inverted splitting tree generation described here.

A Method for Easily Determining Coupling Constant Values: An Addendum to “A Practical Guide to First-Order Multiplet Analysis in ^1H NMR Spectroscopy”

Thomas R. Hoye* and Hongyu Zhao

Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455

hoye@chem.umn.edu

Received July 26, 2000 (Revised Manuscript Received March 28, 2002)

A systematic procedure to decipher first-order ^1H NMR multiplets is described. This method is a very practical tool for extracting coupling constant values. It requires only that one (a) learn to identify each of the 2^n (n = number of spin $1/2$ nuclei to which the proton is coupled) “units of intensity” of a multiplet and (b) then apply a clearly delineated sequence of iterative steps that allows the J s to be assigned in order (from smallest to largest). The approach is even easier to use than one described previously (*J. Org. Chem.* **1994**, *59*, 4096–4103).

Introduction

Several years ago, we described a method for deciphering the individual coupling constants from a complex, first-order multiplet in a ^1H NMR spectrum.¹ Anecdotal evidence makes it clear that many have found this discussion to be very helpful. We now present an alternative, complementary protocol for obtaining the same information. The method we describe here makes the task of extracting the individual J values from complex multiplets even easier.

In general,² a first-order multiplet with chemical shift δ , having coupling constants arising from interaction with n spin $1/2$ nuclei, will contain a maximum of 2^n peaks that are symmetrically arrayed about the midpoint (δ). The actual number of individual peaks is usually reduced by degeneracies arising from certain relationships among two or more J s (e.g., two identical J values or cases where a third J is the sum or difference of two other J values). Nonetheless, the total “units of intensity”, hereafter called components, will always sum to 2^n regardless of the presence or absence of line degeneracy. A trivial example makes the point: a 1:1:1 doublet of doublets (dd) and a 1:2:1 triplet (i.e., a dd with identical J s in which components number 2 and 3 are superimposed) both contain a total of four (2^2) components.³ The increasingly complex set of multiplets shown in Figure 1 reinforce this important point. Recognize that each of the 2^n individual components coincides with one of the following line positions in the multiplet: δ (Hz) + $1/2(\pm J_1 \pm J_2 \pm J_3 \dots \pm J_n)$. The furthest downfield and upfield




Figure 1. Four examples of assignment of all 2^n components to the individual lines in a first-order multiplet.

lines in the multiplet are δ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n)$ and δ (Hz) + $1/2(-J_1 - J_2 - J_3 \dots - J_n)$, respectively. It follows that the separation between the outermost peaks of a multiplet is the sum of the individual coupling constants.^{4,2}

In what follows and for the sake of generality, we consider any multiplet arising from interaction with n protons to be a series of n doublets (e.g., dddd for $n = 4$) regardless of whether any of the protons are chemically and magnetically equivalent. In the case of equivalency,

(4) I.e., the total width of the multiplet = δ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n) - [\delta$ (Hz) + $1/2(-J_1 - J_2 - J_3 \dots - J_n)] = (J_1 + J_2 + J_3 \dots + J_n)$.

(1) Hoye, T. R.; Hanson, P. R.; Vyvyan, J. R. *J. Org. Chem.* **1994**, *59*, 4096–4103.

(2) Most of the concepts noted in this introductory paragraph are well established and accepted truths, the original statements of which are difficult to locate but the existences of which are evidenced in many of the early treatises in the field: e.g., (a) Jackman, L. M.; Sternhell, S. *Applications of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry*; Pergamon: Oxford, 1959. (b) Bovey, F. A. *Nuclear Magnetic Resonance Spectroscopy*; Academic Press: New York, 1969. (c) Becker, E. D. *High-Resolution NMR: Theory and Chemical Applications*; Academic Press: New York, 1969.

(3) The outermost peaks in the multiplet are always the smallest and should be assigned a relative intensity value of 1.

Chart 1. Steps for Identifying J Values in Sequence from Smallest to Largest (J_1 to J_n) ($\{1$ to $x\}$ Represents the Distance between the Peak Corresponding to Component 1 to the Peak Corresponding to Component x)

- i) $\{1$ to $2\}$ is J_1 .^a
- ii) $\{1$ to $3\}$ is J_2 .^b
- iii) remove from further consideration the *component*^c corresponding to $(J_1 + J_2)$.
- iv) $\{1$ to next higher *remaining* component (i.e., 4 or 5) is J_3 . Thus, if $J_1 + J_2$ is $\{1$ to $4\}$, then $\{1$ to $5\}$ is J_3 . Conversely, if $J_1 + J_2$ is not $\{1$ to $4\}$, then $\{1$ to $4\}$ is J_3 .^d
corollary: one of $\{1$ to $4\}$ or $\{1$ to $5\}$ is J_3 .
- v) remove from further consideration the *components*^e corresponding to the remaining combinations of the first three J values [i.e., $(J_1 + J_3)$, $(J_2 + J_3)$, and $(J_1 + J_2 + J_3)$].
- vi) $\{1$ to the *lowest remaining* component} is J_4 .
corollary: one of $\{1$ to $5\}$ through $\{1$ to $9\}$ is J_4 .
- vii) remove from further consideration the *components*^e corresponding to the remaining combinations of the first four J values [i.e., $(J_1 + J_4)$, $(J_2 + J_4)$, $(J_3 + J_4)$, $(J_1 + J_2 + J_4)$, $(J_1 + J_3 + J_4)$, $(J_2 + J_3 + J_4)$, and $(J_1 + J_2 + J_3 + J_4)$].
- viii) $\{1$ to the *lowest remaining* component} is J_5 .
corollary: one of $\{1$ to $6\}$ through $\{1$ to $17\}$ is J_5 .
- ix) etc.

^a $\{1$ to $2\} = \delta$ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n) - [\delta$ (Hz) + $1/2(-J_1 + J_2 + J_3 \dots + J_n)] = J_1$. ^b $\{1$ to $3\} = \delta$ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n) - [\delta$ (Hz) + $1/2(+J_1 - J_2 + J_3 \dots + J_n)] = J_2$. ^c Each peak in the multiplet may (and often will) contain more than one component; each component should be individually removed from consideration or associated with a J value. ^d For a ddd: $\{1$ to $4\}$ (or $\{1$ to $5\}) = \delta$ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n) - [\delta$ (Hz) + $1/2(-J_1 - J_2 + J_3 \dots + J_n)] = J_1 + J_2$ and $\{1$ to $5\}$ (or $\{1$ to $4\}) = \delta$ (Hz) + $1/2(+J_1 + J_2 + J_3 \dots + J_n) - [\delta$ (Hz) + $1/2(+J_1 + J_2 - J_3 \dots + J_n)] = J_3$.




Figure 2. Assignment of J_1 – J_4 for a dddd by applying steps i–vi from Chart 1.

the multiplets are more commonly referred to, of course, as triplets, quartets, pentets, etc. (i.e., dd, ddd, dddd, etc. with identical J s). Recognize then, that the method presented below for deducing the values of all J s in, e.g., a dddd is applicable to *all* multiplets with 16 components: i.e., a pentet (p), a doublet of quartets (dq), a triplet of triplets (tt), or a doublet of doublet of triplets (ddt) in addition to a true doublet of doublet of doublet of doublets (dddd).

The method we now describe for deducing the individual J values from any first-order multiplet requires two principal operations: (i) assignment of each of the individual 2^n components (cf. Figure 1) and (ii) systematic

identification of the individual J s (cf. Chart 1 and the example in Figure 2).⁵

For the first operation, assign every peak in the multiplet one or more component numbers from 1 to 2^n from left to right (arbitrarily) by analogy to the examples shown in Figure 1. This involves assigning the relative intensities among all peaks of the multiplet, an operation that for some complex resonances (cf., Figure 1d) might require an iterative approach but that is usually straightforward for multiplets having 4, 8, 16, and even 32 components.

The second operation requires systematic identification of the J s by the series of steps outlined in Chart 1. Adopt the convention that $J_1 \leq J_2 \leq J_3 \leq J_4 \leq \dots \leq J_n$. Appreciate that for J_3 and beyond it is necessary to have first determined the previous coupling constants (e.g., both J_1 and J_2 must be known before J_3 can be determined). In other words, one must deduce the J s in order, from smallest to largest. Assign individual coupling constant values starting with step i (Chart 1), where $\{1$ to $x\}$ is the distance in Hz between component 1 (which, necessarily, corresponds to the lefthandmost peak) and component x . The task is complete (all J s identified) following step iv for a ddd, vi for a dddd, and viii for a ddddd. In practice, for some complex, slightly non-first-order (leaning/distorted), and/or partially overlapped multiplets, it is advantageous to work synergistically from both extremities of the multiplet.

Consider the dddd, whose component numbers were assigned in Figure 1c. In Figure 2, the determination of

(5) The process requires practice at the outset, but it is well worth learning. Once familiar with the technique, people have deduced the six individual J values in the multiplet shown in Figure 3e (a ddddd) in 1–2 min.




Figure 3. Examples of experimental multiplets for which the complete set of J values has been determined.

the J s, arrived at by sequential application of steps i–vi, is shown.

This method has proven to be very practical. Commonly encountered first-order multiplets can be quickly analyzed.⁶ Some selected multiplets from actual spectra are presented in Figure 3 to further demonstrate the power of this method. Notice that resolution enhancement/line broadening (vertical arrow in Figure 3e) can sometimes make the assignment of the 2nd component numbers more straightforward. Finally, it should not be overlooked that coupling constant values are valuable because they convey information about geometry.^{7,8}

(6) **Note Added in Proof.** For a very recently described, complementary “approach to automated first-order multiplet analysis”, see: Golotvin, S.; Vodopianov, E.; Williams, A. *Magn. Reson. Chem.* **2002**, *40*, 331–336.

(7) Karplus, M. *J. Chem. Phys.* **1959** *30*, 11–15.

Acknowledgment. This study was supported by a grant awarded by the DHHS (CA-76497). We thank Mr. B. M. Eklov for help with rendering the graphics.

JO001139V

(8) For example, the *vicinal* coupling constants for the two protons shown in Figure 3d,e suggest that the hydrogen-bonded conformation **1** (with two gauche relationships between H_a and its methine neighbors and one gauche and two anti arrangements between H_b and its three neighbors) is a major contributor in $CDCl_3$ to the family of conformers that define the solution structure of this polyol fragment. This is not necessarily to be expected because of the steric congestion flanking the C(4)–C(5) bond and would be a very difficult issue to assess by any method other than the *magnitudes* of coupling constants.

